Application Serial No. 10/021,421 Attorney's Docket No. 032751-073

CLAIM SUMMARY DOCUMENT

Claims 1-44 (Canceled)

Claim 45 (Currently Amended) A pharmaceutical preparation comprising an effective amount of at least one complex, wherein said complex comprises:

(i) at least one compound of the formula:

wherein:

the R_a , R_b and R_c residues are, independently of each other, a hydrogen atom or a group of formula II:

with wherein:

 R_1 and R_2 are, independently of each other, C_6 - C_{23} alkyl or alkenyl radicals, which are linear or branched, or radicals -C(=O)- $(C_6$ - $C_{23})$ alkyl or -C(=O)- $(C_6$ - $C_{23})$ alkenyl, which are linear or branched, aryl radicals, cycloalkyl radicals, fluoroalkyl radicals, polyethylene glycol groups, oxyethylene or oxymethylene groups, with the proviso that when one of the R_1 or R_2 is a polyethylene glycol group, an oxyethylene or an oxymethylene group, the other is a linear or branched C_6 - C_{23} alkyl or alkenyl radical, a

linear or branched $-C(=O)-(C_0-C_{22})$ alkyl or $-C(=O)-(C_0-C_{22})$ alkenyl radical , an aryl radical, a cycloalkyl radical, or a fluoroalkyl radical,

p is a positive integer from 1 to 4,

n is a positive integer from 1 to 6,

 m_1 and m_2 are a positive integer from 1 to 6, and $m_1^+ m_1$ may be different for each motif -(CH₂) m_1 -NR₆, and wherein the total number of groups of formula II in said compound is between 1 and 4, wherein said compound is a cationic compound;

- (ii) at least one active substance comprising at least one negative charge; and
- (iii) at least one adjuvant which enhances the formation of the complex between $\int \omega \, C$, said compound and said active substance;

in combination with a pharmaceutically acceptable carrier.

Claim 46 (Previously Added) The preparation of claim 45, comprising, in addition, at least one adjuvant which enhances the introduction of the active substance, comprised in said complex, into a cell.

Claim 47 (Currently Amended) The preparation of claim 46, wherein said adjuvant comprises chloroquine, a protic polar compound chosen from propylene glycol, polyethylene glycol, glycerol, ethanol, 1-methyl-L-2pyrrolidone or derivatives thereof, or